

## COVERINGS BY OPEN CELLS

MÁRIO J. EDMUNDO, PANTELIS E. ELEFThERIOU, AND LUCA PRELLI

ABSTRACT. We prove that in a semi-bounded o-minimal expansion of an ordered group every non-empty open definable set is a finite union of open cells.

## 1. INTRODUCTION

We fix an arbitrary o-minimal expansion  $\mathcal{R} = \langle R, <, +, 0, \dots \rangle$  of an ordered group. Recall that by [3]  $\mathcal{R}$  is semi-bounded if it has no poles; that is, in  $\mathcal{R}$  there is no definable bijection between a bounded and an unbounded interval. See [3] for other characterizations of semi-boundedness. In this note we prove the following theorem.

**Theorem 1.1.** *If  $\mathcal{R}$  is semi-bounded, then every non-empty open definable set is a finite union of open cells.*

As explained in [13, Subsection 2.1], there are three possibilities for an arbitrary o-minimal expansion  $\mathcal{R} = \langle R, <, +, 0, \dots \rangle$  of an ordered group:

- (A)  $\mathcal{R}$  is linear (that is, its first-order theory  $\text{Th}(\mathcal{R})$  is linear ([10])). In this case by [10], there exists  $\mathcal{S} \equiv \mathcal{R}$  with  $\mathcal{S}$  a reduct of an ordered vector space  $\mathcal{V} = \langle V, <, +, 0, \{d\}_{d \in D} \rangle$  over an ordered division ring  $D$  (with the same addition and linear ordering the underlying group of  $\mathcal{S}$ ).
- (B)  $\mathcal{R}$  is not linear. In this case, the theory of every interval in  $\mathcal{R}$  with the induced structure is not linear and so no interval in  $\mathcal{R}$  is elementarily equivalent to a reduct of an interval in an ordered vector space ([10])). Therefore, by the Trichotomy theorem ([14, Theorem 1.2]), a real closed field whose ordering agrees with that of  $\mathcal{R}$  is definable on some interval  $(-e, e)$ . There are now two sub-cases to consider:
  - (B1)  $\mathcal{R}$  is semi-bounded.
  - (B2)  $\mathcal{R}$  is not semi-bounded. In this case, one can endow the whole structure  $\mathcal{R}$  with a definable real closed field. Indeed, let  $\sigma : (a, b) \rightarrow (c, +\infty)$  be a pole in  $\mathcal{R}$ ; that is, a definable bijection (with say,  $\lim_{t \rightarrow b} \sigma(t) = +\infty$ ). Without loss of generality, and using translations, we may assume that  $a = c = 0$  and  $b < e$ . But then, being inside a real closed field, the intervals  $(0, e)$  and  $(0, b)$  are in definable bijection and so  $(0, e)$  and  $(0, +\infty)$  are in definable bijection. Now it is easy to get a real closed field on the whole of  $\mathcal{R}$ .

---

*Date:* March 14, 2013.

*2010 Mathematics Subject Classification.* 03C64.

The first author was supported by Fundação para a Ciência e a Tecnologia, Financiamento Base 2008 - ISFL/1/209. The second author was supported by the Fundação para a Ciência e a Tecnologia grant SFRH/BPD/35000/2007. The third author was supported by Marie Curie grant PIEF-GA-2010-272021. This work is part of the FCT project PTDC/MAT/101740/2008.

*Keywords and phrases:* O-minimal structures, open cells, semi-bounded structures.

A version of Theorem 1.1 in the field case (B2) was proved by Wilkie in [16], for bounded open definable subsets. There are simple examples that show that in this case the boundedness assumption is required. On the other hand, a version of Theorem 1.1 in the linear case (A) was proved by Andrews in [1]. Here we generalize these two results to the semi-bounded non-linear case. Moreover, we also prove a stronger result in the linear case, which we state next. For the notion of ‘linear decomposition’ and ‘star’, see Section 2 below. For the notion of ‘stratification’, see [2, Chapter 4, (1.11)]. By Lemma 2.6, Corollary 2.15 and Proposition 2.17 below, we have:

**Theorem 1.2.** *Assume that  $\mathcal{R} = \langle R, <, 0, +, \{\lambda\}_{\lambda \in D} \rangle$  is an ordered vector space over an ordered division ring  $D$ . Let  $\mathcal{D}$  be a linear decomposition of  $R^n$ . Then there is decomposition  $\mathcal{C}$  of  $R^n$  that refines  $\mathcal{D}$ , such that for every  $C \in \mathcal{C}$ , the star of  $C$  is an open (usual) cell. Moreover,  $\mathcal{C}$  is a stratification of  $R^n$ .*

An important example of a semi-bounded, non-linear o-minimal structure is the expansion  $\mathcal{B}$  of the real ordered vector space  $\mathbb{R}_{\text{vect}} = \langle \mathbb{R}, <, +, 0, \{d\}_{d \in \mathbb{R}} \rangle$  by all bounded semi-algebraic sets. Every bounded interval in  $\mathcal{B}$  admits the structure of a definable real closed field. For example, the field structure on  $(-1, 1)$  induced from  $\mathbb{R}$  via the semi-algebraic bijection  $x \mapsto \frac{x}{\sqrt{1+x^2}}$  is definable in  $\mathcal{B}$ . By [15, 11, 12],  $\mathcal{B}$  is the unique structure that lies strictly between  $\mathbb{R}_{\text{vect}}$  and the real field. The situation becomes significantly more subtle when  $\mathcal{R}$  is non-archimedean, and the study of definable sets and groups in the general semi-bounded setting has recently regained a lot of interest ([4, 6, 7, 8, 13]).

We expect that our main theorem on coverings by open cells (Theorem 1.1) will find numerous applications in the theory of locally definable manifolds in o-minimal structures. Some of those are exhibited in [5]. As stated in that reference, a strengthened result of coverings would yield further applications. We state the desired result here as a Conjecture:

**Conjecture.** Every definable set is a finite union of relatively open definable subsets which are definably simply connected.

*Structure of the paper.* Section 2 contains the stratification result (Theorem 1.2) for the linear case. Section 3 contains the covering by open cells (Theorem 1.1) for the semi-bounded non-linear case.

*Notation.* We recall the standard notation for graphs and “generalized cylinders” of definable maps.

- If  $f : X \rightarrow R$  is a definable map, we denote by  $\Gamma(f)$  the graph of  $f$ .
- If  $f, g : X \rightarrow R$  are definable maps or the constant maps  $-\infty$  and  $+\infty$  on  $X$  with  $f(x) < g(x)$  for all  $x \in X$ , we write  $f < g$  and set:
 
$$(f, g)_X = \{(x, y) \in X \times R : f(x) < y < g(x)\};$$

$$[f, g)_X = \{(x, y) \in X \times R : f(x) \leq y < g(x)\};$$

$$(f, g]_X = \{(x, y) \in X \times R : f(x) < y \leq g(x)\};$$

$$[f, g]_X = \{(x, y) \in X \times R : f(x) \leq y \leq g(x)\}.$$

We also use the same notation for functions  $f, g : Y \rightarrow R$  whose domain  $Y$  contains  $X$  and whose restrictions on  $X$  are as above.

*Acknowledgements.* We wish to thank the referee for many helpful comments.

## 2. THE LINEAR CASE

We assume in this section that  $\mathcal{R} = \langle R, <, 0, +, \{\lambda\}_{\lambda \in D} \rangle$  is an ordered vector space over an ordered division ring  $D$ . For basic properties on such o-minimal structures we refer the reader to [2, Chapter 1, Section 7].

A *linear (affine) function* on  $A \subseteq R^n$  is a function  $f : A \rightarrow R$  of the form  $f(x_1, \dots, x_n) = \lambda_1 x_1 + \dots + \lambda_n x_n + a$ , for some fixed  $\lambda_i \in D$  and  $a \in R$ . For a definable set  $X \subseteq R^n$ , we set  $L(X) = \{f : X \rightarrow R : f \text{ is linear}\}$  and  $L_\infty(X) = L(X) \cup \{\pm\infty\}$ , where we regard  $-\infty$  and  $+\infty$  as constant functions on  $X$ . Then,

- a *linear cell in  $R$*  is either a singleton subset of  $R$ , or an open interval with endpoints in  $R \cup \{\pm\infty\}$ ,
- a *linear cell in  $R^{n+1}$*  is a set of the form  $\Gamma(f)$ , for some  $f \in L(X)$ , or  $(f, g)_X$ , for some  $f, g \in L_\infty(X)$ ,  $f < g$ , where  $X$  is a linear cell in  $R^n$ .

In either case,  $X$  is called *the domain* of the defined cell.

We refer the reader to [2, Chapter 3, (2.10)] for the definition of a *decomposition* of  $R^n$ . A *linear decomposition* of  $R^n$  is then a decomposition  $\mathcal{C}$  of  $R^n$  such that each  $B \in \mathcal{C}$  is a linear cell. The following can be proved similarly to [2, Chapter 3, (2.11)].

**Theorem 2.1** (Linear CDT).

- (1) Given any definable sets  $A_1, \dots, A_k \subseteq R^n$ , there is a linear decomposition  $\mathcal{C}$  of  $R^n$  that partitions each  $A_i$ .
- (2) Given a definable function  $f : A \rightarrow R$ , there is a linear decomposition  $\mathcal{C}$  of  $R^n$  that partitions  $A$  such that the restriction  $f|_B$  to each  $B \in \mathcal{C}$  with  $B \subseteq A$  is linear.

**Definition 2.2.** Let  $\mathcal{C}$  be a linear decomposition of  $R^n$  and  $X$  a subset of  $R^n$ . Denote

$$\text{Star}_{\mathcal{C}}(X) = \{D \in \mathcal{C} : X \cap \text{cl}(D) \neq \emptyset\}.$$

The *star* of  $X$  with respect to  $\mathcal{C}$ , denoted by  $\text{st}_{\mathcal{C}}(X)$ , is then

$$\text{st}_{\mathcal{C}}(X) = \bigcup \text{Star}_{\mathcal{C}}(X).$$

We just write  $\text{Star}(X)$  and  $\text{st}(X)$  if  $\mathcal{C}$  is clear from the context.

In what follows, if  $k > 0$ , then  $\pi : R^{k+1} \rightarrow R^k$  denotes the usual projection map onto the first  $k$ -coordinates, and if  $\mathcal{C}$  is a linear decomposition of  $R^{k+1}$ , then  $\pi(\mathcal{C})$  denotes the linear decomposition  $\{\pi(C) : C \in \mathcal{C}\}$  of  $R^k$ .

**Lemma 2.3.** Let  $\mathcal{C}$  be a linear decomposition of  $R^n$  and  $X$  a subset of  $R^n$ . Then:

- (i) If  $n > 1$ , then  $\text{Star}_{\pi(\mathcal{C})}(\pi(X)) = \pi(\text{Star}_{\mathcal{C}}(X))$ .
- (ii) If  $X$  is an open union of cells in  $\mathcal{C}$ , and  $C \in \mathcal{C}$  with  $C \subseteq X$ , then  $\text{st}(C) \subseteq X$ .

**Proof.** (i)  $\subseteq$ . Let  $D \in \text{Star}(\pi(X))$ . Since  $\pi$  is open, for any open set  $U$  containing  $X$ ,  $\pi(U)$  is an open set containing  $\pi(X)$ . Thus  $D \cap \pi(U) \neq \emptyset$ , which implies  $\pi^{-1}(D) \cap U \neq \emptyset$ . Hence, by the definition of linear decomposition, there is some  $D' \in \text{Star}(X)$  such that  $\pi(D') = D$ .

$\supseteq$ . Let  $D \in \text{Star}(X)$ . For any open set  $U$  containing  $\pi(X)$ ,  $\pi^{-1}(U)$  is an open neighborhood of  $X$ . Therefore  $\pi^{-1}(U) \cap D \neq \emptyset$ , and  $U \cap \pi(D) \neq \emptyset$ . Hence  $\pi(D)$  belongs to  $\text{Star}(\pi(X))$ .

(ii) Since  $X$  is open, for every  $B \in \text{Star}(C)$ ,  $B \cap X \neq \emptyset$ , and hence  $B \subseteq X$ .  $\square$

One would expect that  $\text{st}_C(X)$  is an open set. However, the following example shows that this is not the case.

**Example 2.4.** Consider points  $a_{-1} < a_0 < a_1 < a_2 < a_3$  in  $R$  and let  $\mathcal{C}$  be a linear decomposition of  $R^2$  that contains the following cells:  $(a_{-1}, a_0) \times (a_0, a_2)$ ,  $(a_0, a_1) \times (a_0, a_2)$ ,  $\{a_0\} \times (a_0, a_1)$ ,  $\{a_0\} \times (a_1, a_3)$  and the point  $(a_0, a_1)$ . Then the star of the point  $(a_0, a_1)$  is the union of the above cells, which is not open.

Below we define a special kind of a linear decomposition  $\mathcal{C}$  of  $R^n$  that remedies the above problem. In fact, such a  $\mathcal{C}$  will give us that every  $\text{st}_C(X)$  is an open (usual) cell (see Proposition 2.17 below). From this we obtain the version of Theorem 1.1 for the linear case (see Corollary 2.18 below).

For every  $h \in L(X)$ , where  $X \subseteq R^k$ , and  $h$  of the form  $h(x_1, \dots, x_k) = \lambda_1 x_1 + \dots + \lambda_k x_k + c$ , we define *the extension of  $h$  to  $R^k$*  to be the linear function  $g : R^k \rightarrow R$  with  $g(x_1, \dots, x_k) = \lambda_1 x_1 + \dots + \lambda_k x_k + c$ . We say that  $g$  *extends*  $h$ . In what follows, if  $h \in L(X)$ ,  $X \subseteq R^k$ , and  $c \in \text{cl}(X)$ , we denote  $h(c) := \lim_{t \rightarrow c} h(t)$ , which always exists and is equal to  $g(c)$ , where  $g$  extends  $h$ . In particular, if  $A \subseteq \text{cl}(X)$ , then  $h|_A$  denotes the restriction of  $g$  to  $A$ .

The next definition is by induction on  $n$ .

**Definition 2.5.** A *special linear decomposition* of  $R$  is any linear decomposition of  $R$ . A *special linear decomposition* of  $R^{k+1}$ ,  $k > 0$ , is a linear decomposition  $\mathcal{C}$  of  $R^{k+1}$  with the following two properties:

- Let  $C, C' \in \mathcal{C}$  be two linear cells of the form

$$C = (f, g)_B \text{ and } C' = (f', g')_{B'},$$

where  $B, B' \subseteq R^k$  are disjoint,  $f < g$  in  $L_\infty(B)$  and  $f' < g'$  in  $L_\infty(B')$ . Then, for every  $c \in \text{cl}(B) \cap \text{cl}(B')$ , if  $\pi^{-1}(c) \cap \text{cl}(C) \cap \text{cl}(C')$  is infinite, then

$$\pi^{-1}(c) \cap \text{cl}(C) = \pi^{-1}(c) \cap \text{cl}(C').$$

Equivalently, for every  $c \in \text{cl}(B) \cap \text{cl}(B')$ , if  $\pi^{-1}(c) \cap \text{cl}(C) \cap \text{cl}(C')$  is infinite, then

$$f(c) = f'(c) \text{ and } g(c) = g'(c).$$

- $\pi(\mathcal{C}) = \{\pi(D) : D \in \mathcal{C}\}$  is a special linear decomposition of  $R^k$ .

Before providing the nice consequences of special linear decompositions, we prove that they always exist.

**Lemma 2.6.** *For any linear decomposition  $\mathcal{D}$  of  $R^n$ , there is a special linear decomposition  $\mathcal{C}$  of  $R^n$  that refines  $\mathcal{D}$  (that is, every linear cell in  $\mathcal{D}$  is a union of linear cells in  $\mathcal{C}$ ).*

**Proof.** By induction on  $n$ . For  $n = 1$ , take  $\mathcal{C} = \mathcal{D}$ . Now assume that  $n = k + 1$  and the lemma holds for  $k > 0$ . Let  $\mathcal{D}$  be a linear decomposition of  $R^{k+1}$ . Choose a finite collection  $\mathcal{F}$  of linear maps  $f : R^k \rightarrow R$  such that any linear map that appears in the definition of a linear cell from  $\mathcal{D}$  is a restriction of a map from  $\mathcal{F}$ . Now set

$$\mathcal{G} = \{\Gamma(f) \cap \Gamma(g) : f, g \in \mathcal{F}\} \text{ and } \mathcal{G}' = \{\pi(A) : A \in \mathcal{G}\}.$$

Clearly,  $\mathcal{G}'$  is a finite collection of definable subsets of  $R^k$ . By (Linear CDT and) the inductive hypothesis, there is a special linear decomposition  $\mathcal{C}'$  of  $R^k$  that partitions each  $B \in \mathcal{G}'$ .

**Claim 2.7.** *For every two distinct  $f, g \in \mathcal{F}$ , and  $X \in \mathcal{C}'$ ,*

$$f|_X < g|_X \text{ or } f|_X = g|_X \text{ or } f|_X > g|_X.$$

**Proof.** Indeed, let  $A = \Gamma(f) \cap \Gamma(g) \neq \emptyset$ . Since  $\pi(A) \in \mathcal{G}'$  and  $\mathcal{C}'$  partitions  $\pi(A)$ , we have either  $X \subseteq \pi(A)$  or  $X \subseteq R^k \setminus \pi(A)$ . If  $X \subseteq \pi(A)$ , then of course  $f|_X = g|_X$ . Suppose that  $X \subseteq R^k \setminus \pi(A)$ , then  $\Gamma(f|_X) \cap \Gamma(g|_X) = \emptyset$ . So for every  $x \in X$  either  $f|_X(x) < g|_X(x)$  or  $g|_X(x) < f|_X(x)$  and therefore  $X = X^- \sqcup X^+$  where  $X^- = \{x \in X : f|_X(x) < g|_X(x)\}$  and  $X^+ = \{x \in X : g|_X(x) < f|_X(x)\}$ . Since both  $X^-$  and  $X^+$  are open definable subsets and  $X$  is definably connected, either  $X = X^-$  or  $X = X^+$ .  $\square$

We can thus write  $\mathcal{C}' = \{X_1, \dots, X_m\}$ , such that for each  $i \in \{1, \dots, m\}$ ,  $f_{i1} < \dots < f_{in(i)}$  are the distinct functions in  $L(X_i)$ , each being a restriction of some  $f \in \mathcal{F}$ , and exhausting all possible such. Then

$$\mathcal{C}_i = \{(-\infty, f_{1|X_i}), (f_{1|X_i}, f_{2|X_i}), \dots, (f_{l|X_i}, \infty), \Gamma(f_{1|X_i}), \dots, \Gamma(f_{l|X_i})\}$$

is a partition of  $\pi^{-1}(X_i)$ , and  $\mathcal{C} = \mathcal{C}_1 \cup \dots \cup \mathcal{C}_k$  is a linear decomposition of  $R^{k+1}$  which refines  $\mathcal{D}$ . We show that  $\mathcal{C}$  is special. Let  $C = (f, g)_B$  and  $C' = (f', g')_{B'}$  be as in Definition 2.5. We need to check that

$$f(c) = f'(c) \text{ and } g(c) = g'(c)$$

for every  $c \in cl(B) \cap cl(B')$ . If not, then since  $\pi^{-1}(c) \cap cl(C) \cap cl(C')$  is infinite, we have either

$$f(c) < f'(c) < g(c) \text{ or } f'(c) < g(c) < g'(c)$$

or

$$f'(c) < f(c) < g'(c) \text{ or } f(c) < g'(c) < g(c).$$

In the first case, the extension  $h$  of  $f'$  restricted to  $B$  satisfies:

$$f < h|_B < g.$$

This contradicts the definition of  $\mathcal{C}_i$  with  $i$  such that  $X_i = B$ . In the second case we also get a contradiction by considering  $i$  such that  $X_i = B'$ . For the remaining two cases the arguments are similar.

Finally, notice that  $\pi(\mathcal{C}) = \mathcal{C}'$  is a special linear decomposition of  $R^k$ .  $\square$

We now aim towards Proposition 2.17 below. We begin with the following lemma. For  $X \subseteq R^{n+1}$  and  $A \subseteq R^n$ , let us write  $X_A = X \cap \pi^{-1}(A)$ .

**Lemma 2.8.** *Let  $C$  be a linear cell with domain  $B$ , and let  $A \subseteq cl(B)$ . Then the following hold:*

- If  $C = (f, g)_B$ , then  $cl(C) = [\bar{f}, \bar{g}]_{cl(B)}$  where  $\bar{f} = f|_{cl(B)}$  and  $\bar{g} = g|_{cl(B)}$ .  
Moreover,  $cl(C)_A = [\bar{f}, \bar{g}]_A$ .
- If  $C = \Gamma(h)$  with  $h : B \rightarrow R$ , then  $cl(C) = \Gamma(\bar{h})$  where  $\bar{h} = h|_{cl(B)}$ .  
Moreover,  $cl(C)_A = \Gamma(\bar{h})_A$ .

**Proof.** We do the case  $C = (f, g)_B$ , since the other case is easier. Let  $c = (b, y) \in cl(C)$  with  $b = \pi(c)$  and  $y \in R$ . Then by curve selection [2, Chapter 6, (1.5)] there is a continuous definable map  $\gamma : (0, \epsilon) \rightarrow C$  such  $\gamma(0) := \lim_{t \rightarrow 0^+} \gamma(t) = c$ . Let  $\delta = \pi \circ \gamma : (0, \epsilon) \rightarrow B$  and let  $\tau : (0, \epsilon) \rightarrow R$  such that  $\gamma(t) = (\delta(t), \tau(t))$ . Since for every  $t$ , we have  $f(\delta(t)) < \tau(t) < g(\delta(t))$ , taking the limit  $\lim_{t \rightarrow 0^+}$  we see that  $f(b) \leq y \leq g(b)$ . Therefore  $c \in [\bar{f}, \bar{g}]_{cl(B)}$  since  $b = \lim_{t \rightarrow 0^+} \delta(t) \in cl(B)$ . So we have  $cl(C) \subseteq [\bar{f}, \bar{g}]_{cl(B)}$ .

Let now  $c = (b, y) \in [\bar{f}, \bar{g}]_{cl(B)}$ . If  $c \in \Gamma(\bar{f})$  (respectively,  $c \in \Gamma(\bar{g})$ ) then any open definable neighborhood of  $c$  intersects  $\Gamma(f)$  (respectively,  $\Gamma(g)$ ) which is contained in  $cl(C)$ . So  $c \in cl(C)$  in this case. Otherwise,  $b \in cl(B)$  and  $\bar{f}(b) < y < \bar{g}(b)$ . Then by curve selection [2, Chapter 6, (1.5)] there is a continuous definable map  $\delta : (0, \epsilon) \rightarrow B$  such  $\delta(0) := \lim_{t \rightarrow 0^+} \delta(t) = b$ . Let  $\gamma : (0, \epsilon) \rightarrow R^n$  be given by  $\gamma(t) = (\delta(t), y)$ . Then by continuity and replacing  $\epsilon$  if necessary we have that  $\gamma((0, \epsilon)) \subseteq C$  and clearly  $\gamma(0) := \lim_{t \rightarrow 0^+} \gamma(t) = c$ . So  $c \in cl(C)$  in this case also. Therefore,  $[\bar{f}, \bar{g}]_{cl(B)} \subseteq cl(C)$ .

For the moreover clause,  $cl(C)_A = cl(C) \cap \pi^{-1}(A) = [\bar{f}, \bar{g}]_{cl(B)} \cap \pi^{-1}(A) = [\bar{f}, \bar{g}]_A$ .  $\square$

We will often use without mentioning the following result, which is proved in arbitrary o-minimal expansions of ordered groups for the case of bounded cells in [2, Chapter 6, (1.7)]. It is remarked there that the boundedness assumption is necessary. However, in the linear case, it is not, and the result follows immediately from the previous lemma.

**Lemma 2.9.** *For any linear cell  $C \subseteq R^n$ ,  $\pi(cl(C)) = cl(\pi(C))$ .*

The following general result about linear cells is required below. It is perhaps already known in the literature, yet we include a (somewhat sketchy) proof for completeness.

**Lemma 2.10.** *Let  $C, D \subseteq R^n$  be two linear cells. Then  $cl(C) \cap cl(D)$  is definably connected.*

*Proof.* We may assume that  $cl(C) \cap cl(D)$  is non-empty. We work by induction on  $n$ . For  $n = 1$ , each of  $C$  and  $D$  is an open interval or a point, and, hence,  $cl(C) \cap cl(D)$  is a closed interval or a point.

Assume now that  $C, D \subseteq R^{n+1}$  and that we know the claim for all pairs of linear cells in  $R^n$ . Let  $A = cl(\pi(C)) \cap cl(\pi(D))$ . By inductive hypothesis,  $A$  is definably connected. We claim that

$$cl(C) \cap cl(D) = cl(C)_A \cap cl(D)_A$$

Indeed, if  $(x, y) \in cl(C) \cap cl(D)$ , then  $x \in \pi(cl(C) \cap cl(D)) \subseteq \pi(cl(C)) \cap \pi(cl(D)) = cl(\pi(C)) \cap cl(\pi(D)) = A$ , by Lemma 2.9.

In what follows, we handle the case where  $C$  and  $D$  are both cylinders (the rest of the cases are easier). Let  $C = (f_1, g_1)_{\pi(C)}$  and  $D = (f_2, g_2)_{\pi(D)}$ . By Lemma 2.8, we have that  $cl(C)_A = [\bar{f}_1, \bar{g}_1]_A$  and  $cl(D)_A = [\bar{f}_2, \bar{g}_2]_A$ , where  $\bar{f}_i = f_i|_A$  and  $\bar{g}_i = g_i|_A$ .

If the graphs of  $\bar{f}_1, \bar{g}_1$  do not intersect those of  $\bar{f}_2, \bar{g}_2$ , then  $cl(C)_A \cap cl(D)_A$  is empty or of the form  $[f, g]_A$ , for some linear maps  $f, g$ .

If the graphs of  $\overline{f}_1, \overline{g}_1$  intersect those of  $f_2, g_2$ , then, by linearity of these functions, it is easy to see that  $cl(C)_A \cap cl(D)_A$  is of the form  $[f, g]_{A'}$ , where

- $A' \subseteq A$  is definably connected, and
- $f$  and  $g$  are piecewise linear and continuous on  $A'$ .

In all these cases, it is easy to check that  $cl(C)_A \cap cl(D)_A$  is definably connected. Indeed, it is a cylinder between two definably connected sets (the graphs of  $f$  and  $g$  above  $A$  or  $A'$ ).  $\square$

Before we proceed we make the following remark.

**Remark 2.11.** Let  $A \subseteq R^n$  be a subset. We say that  $A$  is *convex* if for all  $x, y \in A$  and for all  $q \in \mathbb{Q} \cap [0, 1]$  we have  $qx + (1 - q)y \in A$ . See [9, Definition 3.1].

The following hold:

- The intersection of two convex sets is convex.
- Every linear cell is convex.
- If  $B$  is a convex definable set, then  $cl(B)$  is also convex.

The first two points are clear. We need to argue that if  $B$  is convex then so is  $cl(B)$ . Let  $x, y \in cl(B)$ , and  $q \in \mathbb{Q} \cap [0, 1]$ . By curve selection [2, Chapter 6, (1.5)] let  $\gamma : (0, \epsilon) \rightarrow B$  and  $\delta : (0, \epsilon) \rightarrow B$  be continuous definable maps with  $\lim_{t \rightarrow 0^+} \gamma(t) = x$  and  $\lim_{t \rightarrow 0^+} \delta(t) = y$ . By convexity of  $B$  we have a continuous definable map  $\tau : (0, \epsilon) \rightarrow B : t \mapsto q\gamma(t) + (1 - q)\delta(t)$ . Now we have  $\lim_{t \rightarrow 0^+} \tau(t) = qx + (1 - q)y \in cl(B)$  as required.

**Lemma 2.12.** Let  $\mathcal{C}$  be a special linear decomposition of  $R^n$ ,  $n > 1$ ,  $D, E \in \mathcal{C}$  two linear cells of the form

$$D = \Gamma(f) \text{ and } E = \Gamma(g),$$

where  $f \in L(B)$ ,  $g \in L(B')$ , and  $A = cl(B) \cap cl(B') \neq \emptyset$ . Then:

$$f|_A < g|_A \text{ or } f|_A = g|_A \text{ or } f|_A > g|_A.$$

**Proof.** Assume not. First we prove the following claim:


**Claim 2.13.** There are points  $c, d \in A$ , such that  $f(c) = g(c)$  and  $f(d) \neq g(d)$ .

**Proof.** Since  $f|_A \neq g|_A$ , then there exists  $d \in A$  such that  $f(d) \neq g(d)$ . On the other hand, by Lemma 2.10,  $A = cl(B) \cap cl(B')$  is definably connected. Therefore  $A$  is definably path connected ([2, Chapter 6, (3.2)]). So now let  $d_1, d_2 \in A$  so that  $f(d_1) < g(d_1)$  and  $f(d_2) > g(d_2)$ . Let  $\gamma : [0, \epsilon] \rightarrow A$  be a path from  $d_1$  to  $d_2$  inside  $A$ , and consider the definable function  $F = (f - g) \circ \gamma : [0, \epsilon] \rightarrow R$ . By the o-minimal intermediate value theorem,  $F$  obtains value 0 at some point  $t_0$  in  $[0, \epsilon]$ . So if  $c = \gamma(t_0) \in A$ , then  $f(c) = g(c)$ .  $\square$

By the claim there are points  $c, d \in A$ , such that  $f(c) = g(c)$  and  $f(d) \neq g(d)$ . Say,  $f(d) < g(d)$ . Let  $F, G \in \mathcal{C}$  be linear cells of the form  $F = (h, k)_B$ ,  $G = (l, m)_{B'}$  such that

$$f|_A = h|_A < k|_A \text{ and } g|_A = l|_A < m|_A.$$

We next claim that there is a point  $e \in A$ , such that  $f(e) < g(e) < k(e)$ .



If  $g(d) < k(d)$ , then let  $e = d$ . So assume  $k(d) \leq g(d)$ . We will choose  $e$  to be “between”  $c$  and  $d$ . We first see that there is  $q_0 \in (0, 1] \cap \mathbb{Q}$ , such that

$$q_0 g(d) + (1 - q_0)g(c) < q_0 k(d) + (1 - q_0)k(c)$$

Indeed, if not, then  $k(c) \leq g(c)$ . But  $g(c) = f(c) < k(c)$ , a contradiction. On the other hand, since  $f(d) < g(d)$  and  $f(c) = g(c)$ , we have that for every  $q \in (0, 1] \cap \mathbb{Q}$ ,

$$q f(d) + (1 - q)f(c) < q g(d) + (1 - q)g(c).$$

Hence, if we let  $e = q_0 d + (1 - q_0)c$ , then  $e \in A$  (by Remark 2.11) and we have  $f(e) < g(e) < k(e)$ , proving our claim.

Now, since  $f(e) = h(e)$  and  $g(e) = l(e)$ , we have  $h(e) < l(e) < k(e)$ . This implies that  $\pi^{-1}(e) \cap cl(F) \cap cl(G)$  is infinite, but  $\pi^{-1}(e) \cap cl(F) \neq \pi^{-1}(e) \cap cl(G)$ , contradicting the fact that  $\mathcal{C}$  is special.  $\square$

**Lemma 2.14.** *Let  $\mathcal{C}$  be a special linear decomposition of  $R^n$ ,  $n > 1$ , and  $D, E \in \mathcal{C}$  such that  $D \cap cl(E) \neq \emptyset$ . Then:*

$$\pi(D) \subseteq cl(\pi(E)) \Rightarrow D \subseteq cl(E).$$

**Proof.** The statement trivially holds if  $D = E$ , hence assume  $D \neq E$ . Let  $E = (f, g)_B$  or  $E = \Gamma(f)$ , for some  $f, g \in L_\infty(B)$ . If  $D$  has domain  $B$ , then  $E = (f, g)_B$ , and  $D = \Gamma(f)$  or  $D = \Gamma(g)$ . Hence,  $D \subseteq cl(E)$ . So now assume that  $D$  has domain  $B' = \pi(D)$ , disjoint from  $B = \pi(E)$ , and, for a contradiction, that  $B' \subseteq cl(B)$  but  $D \not\subseteq cl(E)$ . Let  $A = cl(B) \cap cl(B') \neq \emptyset$ .

Case A:  $D = \Gamma(g')$ , for some  $g' \in L(B')$ . Then one of the pairs  $f, g'$  or  $g, g'$  must contradict Lemma 2.12.

Case B:  $D = (f', g')_{B'}$ , for some  $f', g' \in L_\infty(B)$ . Then, again by Lemma 2.12, applied to each of the four pairs  $\{f, f'\}$ ,  $\{f, g'\}$ ,  $\{g, f'\}$ ,  $\{g, g'\}$  that are involved, the only remaining possibilities are the following:

$$f'_{|A} < g_{|A} < g'_{|A} \text{ or } f'_{|A} < f_{|A} < g'_{|A}.$$

In the first case, let  $F \in \mathcal{C}$  be a linear cell of the form  $F = (h, k)_B$ , such that

$$g_{|A} = h_{|A} < k_{|A}.$$

Then for any  $c \in A$ ,  $f'(c) < h(c) < g'(c)$ . This implies that  $\pi^{-1}(c) \cap cl(D) \cap cl(F)$  is infinite, but  $\pi^{-1}(c) \cap cl(D) \neq \pi^{-1}(c) \cap cl(F)$ , contradicting the fact that  $\mathcal{C}$  is special. Similarly for the second case.  $\square$

**Corollary 2.15.** *Let  $\mathcal{C}$  be a special linear decomposition of  $R^n$ ,  $n > 0$ , and  $D, E \in \mathcal{C}$  such that  $D \cap cl(E) \neq \emptyset$ . Then  $D \subseteq cl(E)$ .*

*In particular,  $\mathcal{C}$  is a stratification of  $R^n$ .*



**Proof.** The statement trivially holds if  $D = E$ , hence assume  $D \neq E$ . We work by induction on  $n$ . For  $n = 1$ , the assumption  $D \cap cl(E) \neq \emptyset$  implies that  $E$  is an open interval and  $D$  is one of its endpoints. So now assume  $n > 1$ . Clearly,  $\pi(D) \cap cl(\pi(E)) \neq \emptyset$  (using Lemma 2.9), and hence by inductive hypothesis,  $\pi(D) \subseteq cl(\pi(E))$ . By Lemma 2.14,  $D \subseteq cl(E)$ .  $\square$

**Lemma 2.16.** *Let  $\mathcal{C}$  be a special linear decomposition of  $R^n$ ,  $n > 0$ . Then, for any subset  $X \subseteq R^n$ ,  $st(X)$  is open.*

**Proof.** It suffices to show that  $st(X) \cap cl(E) = \emptyset$  for any  $E \in \mathcal{C}$  with  $st(X) \cap E = \emptyset$ . Suppose this is not the case. Then some  $D \in \text{Star}(X)$  meets  $cl(E)$ . Then by Corollary 2.15,  $cl(E)$  contains  $D$  and so  $cl(D)$ . As  $X$  meets  $cl(D)$ , it meets  $cl(E)$ , and hence  $E \subseteq st(X)$ , which is a contradiction.  $\square$

**Proposition 2.17.** *Let  $\mathcal{C}$  be a special linear decomposition of  $R^n$ ,  $n > 0$ , and  $C \in \mathcal{C}$ . Then  $U = st(C)$  is an open (usual) cell.*

**Proof.** By Lemma 2.16,  $U$  is open and thus has dimension  $n$ . If  $\dim(C) = n$ , then  $U = C$  and the statement holds trivially. We may thus assume that  $\dim(C) < n$ . We work by induction on  $n$ . If  $n = 1$ , then  $C$  is a point and  $U$  is an open interval. Now assume that  $n = k + 1$  and the Claim holds for  $k > 0$ .

Assume first that  $C$  is a linear cell in  $\mathcal{C}$  of dimension  $k$  which is the graph of a linear function  $h : D \rightarrow R$ . Then clearly

$$st(C) = (f, g)_D,$$

for some  $f, g \in L_\infty(D)$  with  $f < h < g$ .

In all other cases,  $\dim(\pi(C)) < \dim(\pi(U))$ . Since  $\mathcal{C}$  is a linear decomposition, for every  $B \in \text{Star}(\pi(C))$ ,  $\pi^{-1}(B) \cap U$  is a union of linear cells in  $\mathcal{C}$  which are either graphs of linear maps, or cylinders between linear maps, with domain  $B$ . By Lemma 2.3(i),  $U \subseteq \bigcup \{\pi^{-1}(B) : B \in \text{Star}(\pi(C))\}$ , and hence

$$U = \bigcup \{\pi^{-1}(B) \cap U : B \in \text{Star}(\pi(C))\}.$$

We claim that for every  $B \in \text{Star}(\pi(C))$ ,

$$\pi^{-1}(B) \cap U = (f_B, g_B)_B,$$

for some  $f_B, g_B \in L_\infty(B)$  with  $f_B < g_B$ .

Fix  $B \in \text{Star}(\pi(C))$ . Let  $f_B$  be the bottom function with domain  $B$  defining the bottom cell of  $\pi^{-1}(B) \cap U$  and let  $g_B$  be the top function with domain  $B$  defining the top cell of  $\pi^{-1}(B) \cap U$ . (Recall that this latter set is a union of linear cells in  $\mathcal{C}$  which are either graphs of linear maps, or cylinders between linear maps, with domain  $B$ ). Let  $y \in \pi(C) \cap cl(B)$ . Since  $C$  is a cell and since the cell in  $\mathcal{C}$  contained in  $\pi^{-1}(B) \cap U$  in which  $f_B$  (respectively,  $g_B$ ) appears intersects  $C$ , we have that  $\pi^{-1}(y) \cap U$  is a cell of the form  $\{y\} \times (f_B(y), g_B(y))$ . But then for every cell  $A \in \mathcal{C}$  with  $\pi(A) = B$  and  $A \subseteq (f_B, g_B)_B$ , the closure  $cl(A)$  must contain a point  $x \in \pi^{-1}(y) \cap U \subseteq C$ . So  $cl(A)$  has non-empty intersection with  $C$ . Since  $U = st(C)$ ,  $A$  is contained in  $U$ . Since  $U$  is open (Lemma 2.16), we easily obtain that for every  $B \in \text{Star}(\pi(C))$ ,

$$\pi^{-1}(B) \cap U = (f_B, g_B)_B,$$

for some  $f_B, g_B \in L_\infty(B)$  with  $f_B < g_B$  as required.

Let  $D = \text{st}(\pi(C))$ ,  $f = \bigcup_{B \in \text{Star}(\pi(C))} f_B$  and  $g = \bigcup_{B \in \text{Star}(\pi(C))} g_B$ . Then

$$U = (f, g)_D.$$

By inductive hypothesis,  $D$  is a usual cell. To show that  $f, g$  are continuous, we need to show that for every  $B, B' \in \text{Star}(\pi(C))$ , and  $c \in \text{cl}(B) \cap \text{cl}(B')$ ,

$$f_B(c) = f_{B'}(c) \text{ and } g_B(c) = g_{B'}(c).$$

Let  $H = (h, g_B)_B$  be the upper-most linear cell in  $\mathcal{C}$  contained in  $(f_B, g_B)_B$  and  $H' = (h', g_{B'})_{B'}$  the upper-most linear cell in  $\mathcal{C}$  contained in  $(f_{B'}, g_{B'})_{B'}$ . By Corollary 2.15,  $C \subseteq \text{cl}(H) \cap \text{cl}(H')$ . Hence, if  $C = (l, m)_A$ , for some  $l, m \in L(A)$ , then  $\pi^{-1}(c) \cap \text{cl}(H) \cap \text{cl}(H')$  is infinite. On the other hand, if  $C = \Gamma(l)$  for some  $l \in L(A)$ , then by Lemma 2.12,

$$h|_A \leq l \text{ and } h'|_A \leq l,$$

and hence  $\pi^{-1}(c) \cap \text{cl}(H) \cap \text{cl}(H')$  is again infinite. Since  $\mathcal{C}$  is special,

$$h(c) = h'(c) \text{ and } g_B(c) = g_{B'}(c).$$

Similarly, we can show that  $f_B(c) = f_{B'}(c)$ .

It follows that  $U = (f, g)_D$  is a cell.  $\square$

**Corollary 2.18.** *If  $\mathcal{R} = (R, <, 0, +, \{\lambda\}_{\lambda \in D})$  is an ordered vector space over an ordered division ring  $D$ , then every non-empty open definable set is a finite union of open cells.*

**Proof.** Let  $X \subseteq R^n$  be an open definable subset and take  $\mathcal{C}$  a special linear decomposition of  $R^n$  that partitions  $X$ . By Lemma 2.3(ii),

$$X = \bigcup_{C \in \mathcal{C}, C \subseteq X} \text{st}(C).$$

Then apply Proposition 2.17.  $\square$

### 3. THE SEMI-BOUNDED NON-LINEAR CASE

We assume in this section that  $\mathcal{R}$  is semi-bounded and non-linear. So, as we saw in the Introduction, there exists a definable real closed field  $\langle I, 0_I, 1_I, +_I, \cdot_I, <_I \rangle$  on some interval  $I \subseteq R$  which, without loss of generality, can be assumed to be of the form  $I = (-e, e)$ ,  $0_I = 0$  and  $<_I$  is the restriction of  $<$  to  $I$ . Here we will use the existence of this “short” definable real closed field to adapt Wilkie’s proof ([16]) in o-minimal expansions of real closed fields.

In the next lemmas the semi-boundedness assumption of  $\mathcal{R}$  is not required.

**Lemma 3.1** ([16], Lemma 1). *Let  $C$  be a cell in  $R^n$ . Then there exists an open cell  $D$  in  $R^n$  with  $C \subseteq D$  and a definable retraction  $H : D \rightarrow C$  (that is, a continuous map such that  $H|_C = \text{id}_C$ ).*

**Lemma 3.2.** *Let  $C$  be a cell in  $R^n$ . Suppose that  $h : C \rightarrow R$  is a continuous definable map and let  $U$  be an open definable subset of  $R^{n+1}$ . Suppose further that  $\Gamma(h) \subseteq U$ . Then there exist definable maps  $f, g : C \rightarrow R$  and cells  $C_1, \dots, C_m \subseteq C$  such that:*

- (1)  $f < h < g$ ;
- (2)  $C = C_1 \cup \dots \cup C_m$ ;

- (3) for each  $i$ ,  $f|_{C_i}$  and  $g|_{C_i}$  are continuous;
- (4) for each  $i$ ,  $\Gamma(h|_{C_i}) \subseteq [f|_{C_i}, g|_{C_i}]_{C_i} \subseteq U$ .

**Proof.** Since  $U$  is open and  $\Gamma(h) \subseteq U$ , by definable choice ([2, Chapter 6, (1.2)]) there exists definable maps  $f, g : C \rightarrow R$  such that  $f < h < g$  and  $[f, g]_C \subseteq U$ . By cell decomposition, there are cells  $C_1, \dots, C_m \subseteq C$  covering  $C$  such that for each  $i$ ,  $f|_{C_i}$  and  $g|_{C_i}$  are continuous. Now the rest is clear.  $\square$

The following is also needed:

**Lemma 3.3.** *Let  $C$  be a cell in  $R^n$ . Suppose that  $f, g : C \rightarrow R$  are continuous definable maps such that  $f < g$  and let  $V, W \subseteq U$  be open definable subsets of  $R^{n+1}$ . Suppose further that  $(f, g)_C \subseteq U$ ,  $\Gamma(f) \subseteq V$  and  $\Gamma(g) \subseteq W$ . Then there exist definable maps  $f', g' : C \rightarrow R$  and cells  $C_1, \dots, C_m \subseteq C$  such that:*

- (1)  $C = C_1 \cup \dots \cup C_m$ ;
- (2) for each  $i$ ,  $f'|_{C_i}$  and  $g'|_{C_i}$  are continuous;
- (3)  $f < f' < g' < g$ ;
- (4) for each  $i$ ,  $\Gamma(f'|_{C_i}) \subseteq V$  and  $\Gamma(g'|_{C_i}) \subseteq W$ ;
- (5) for each  $i$ ,  $(f'|_{C_i}, g'|_{C_i})_{C_i} \subseteq U$ ,  $(f|_{C_i}, g|_{C_i})_{C_i} \subseteq U$  and  $[f'|_{C_i}, g'|_{C_i}]_{C_i} \subseteq U$ .

**Proof.** Since  $(f, g)_C \subseteq U$ ,  $\Gamma(f) \subseteq V$  and  $\Gamma(g) \subseteq W$  and  $V, W \subseteq U$  be open definable subsets of  $R^{n+1}$ , by definable choice ([2, Chapter 6, (1.2)]) there exists definable maps  $f', g' : C \rightarrow R$  such that

- (1)  $f < f' < g' < g$ ;
- (2)  $\Gamma(f') \subseteq V$  and  $\Gamma(g') \subseteq W$ ;
- (3)  $(f', g')_C \subseteq U$ ,  $(f, g)_C \subseteq U$  and  $[f', g']_C \subseteq U$ .

By cell decomposition, there are cells  $C_1, \dots, C_m \subseteq C$  covering  $C$  such that for each  $i$ ,  $f|_{C_i}$  and  $g|_{C_i}$  are continuous. Now the rest is clear.  $\square$

Below we let

$$d^{(n)}(\bar{x}, \bar{y}) = \max\{|x_i - y_i| : 1 \leq i \leq n\}$$

denote the standard distance in  $R^n$  (where we denote by  $\bar{z} = (z_1, \dots, z_n)$  the elements of  $R^n$ ). This distance is a continuous definable function (by [2, Chapter 6 (1.4)]). Moreover, if  $B \subseteq R^n$  is a nonempty definable subset and  $\bar{a} \in R^n$ , then

$$d^n(\bar{a}, B) = \inf\{d^n(\bar{a}, \bar{x}) : \bar{x} \in B\}$$

is well defined (by ([2, Chapter 1 (3.3)])) and  $d^n(\bar{a}, B) = 0$  if and only if  $\bar{a} \in cl(B)$  (the if part of this equivalence is immediate and for the only if part one can use the curve selection ([2, Chapter 6 (1.5)])).

Let  $\pi : R^{n+1} \rightarrow R^n$  be the projection onto the first  $n$  coordinates. We say that an open definable subset  $U$  of  $R^{n+1}$  has *I-short height* if for every  $\bar{x} \in \pi(U)$  we have

$$\sup\{|t - s| : t, s \in U_{\bar{x}}\} \in I$$

where  $U_{\bar{x}} = \{y \in R : (\bar{x}, y) \in U\}$ .

We now prove the analogue of [16, Lemma 2] for open definable subsets with *I-short height*. The argument of the proof is similar, one just has to observe that the field operations are used in Wilkie's proof in a uniform way and only along

fibers. Since in our case our fibers are  $I$ -short, such field operations, in the field  $I$ , can also be used in exactly the same way.

For completeness we include the details of the proof but at the end we follow a more constructive argument suggested to us by Oleg Belagradek. For that we need the following observations which are true in arbitrary o-minimal expansions of ordered groups:

**Remark 3.4.** If  $\theta : [a, b] \rightarrow [c, d]$  is a continuous, definable, strictly decreasing function,  $\theta(a) = d$  and  $\theta(b) = c$ , then  $\theta$  is bijective.

Indeed, as  $\theta$  is definable and continuous,  $\theta([a, b])$  is definable, closed, and bounded by [2, Chapter 6 (1.10)], and hence it is a finite union of closed intervals and singletons, by o-minimality. Since  $\theta$  is strictly decreasing,  $\theta([a, b])$  is densely ordered, and so is a closed interval, which must be  $[c, d]$ .

**Remark 3.5.** Let  $V \subseteq R^n$  be an open definable subset, and let  $\{\theta_{\bar{x}} : [a_{\bar{x}}, b_{\bar{x}}] \rightarrow [c_{\bar{x}}, d_{\bar{x}}]\}_{\bar{x} \in V}$  be a uniformly definable family of strictly decreasing functions with  $\theta_{\bar{x}}(a_{\bar{x}}) = d_{\bar{x}}$  and  $\theta_{\bar{x}}(b_{\bar{x}}) = c_{\bar{x}}$ . (So by the previous remark all  $\theta_{\bar{x}}$ 's are bijective). Suppose that all  $a_{\bar{x}}, b_{\bar{x}}, c_{\bar{x}}, d_{\bar{x}}$  are continuous functions in  $\bar{x}$ , and moreover, the map

$$\{(\bar{x}, y) : \bar{x} \in V \text{ and } y \in [a_{\bar{x}}, b_{\bar{x}}]\} \rightarrow R : (\bar{x}, y) \mapsto \theta_{\bar{x}}(y)$$

is continuous. Then the map

$$\gamma : \{(\bar{x}, z) : \bar{x} \in V \text{ and } z \in [c_{\bar{x}}, d_{\bar{x}}]\} \rightarrow R : (\bar{x}, y) \mapsto \theta_{\bar{x}}^{-1}(z)$$

is continuous.

Indeed, for each  $\bar{x}$ , let  $\bar{\theta}_{\bar{x}} : R \rightarrow R$  be given by

$$\bar{\theta}_{\bar{x}}(y) = \begin{cases} a_{\bar{x}} + d_{\bar{x}} - y & \text{for } y < a_{\bar{x}} \\ \theta_{\bar{x}}(y) & \text{for } a_{\bar{x}} \leq y \leq b_{\bar{x}} \\ b_{\bar{x}} + c_{\bar{x}} - y & \text{for } y > b_{\bar{x}} \end{cases}$$

Then  $\{\bar{\theta}_{\bar{x}} : R \rightarrow R\}_{\bar{x} \in V}$  is a uniformly definable family of strictly decreasing functions such that  $\bar{\theta}_{\bar{x}}$  extends  $\theta_{\bar{x}}$  for all  $\bar{x} \in V$ , and  $V \times R \rightarrow R : (\bar{x}, y) \mapsto \bar{\theta}_{\bar{x}}(y)$  is a continuous function.

Now  $\bar{\gamma} : V \times R \rightarrow R : (\bar{x}, y) \mapsto \bar{\theta}_{\bar{x}}^{-1}(z)$  is also a continuous function since, for any  $(a, b) \subseteq R$ ,

$$\begin{aligned} \bar{\gamma}^{-1}((a, b)) &= \{(\bar{x}, z) \in V \times R : a < \bar{\theta}_{\bar{x}}^{-1}(z) < b\} \\ &= \{(\bar{x}, z) \in V \times R : \bar{\theta}_{\bar{x}}(a) < z < \bar{\theta}_{\bar{x}}(b)\}, \end{aligned}$$

which is open. Therefore, since  $\bar{\gamma}$  extends  $\gamma$ , we have that  $\gamma$  is also continuous as required.

**Lemma 3.6.** Let  $C$  be a cell in  $R^n$ . Suppose that  $f, g : C \rightarrow R$  are continuous definable maps such that  $f < g$  and let  $U$  be an open definable subset of  $R^{n+1}$  with  $I$ -short height. Suppose further that  $[f, g]_C \subseteq U$  (respectively  $(f, g)_C \subseteq U$ ). Then there exists an open definable subset  $V$  of  $R^n$  and continuous definable maps  $F, G : V \rightarrow R$  such that:

- (1)  $C \subseteq V$ ;
- (2)  $F|_C = f$  and  $\Gamma(F) \subseteq U$  (respectively  $\Gamma(G) \subseteq U$ );
- (3)  $G|_C = g$ ;
- (4)  $F < G$ ;

- (5) for all  $\bar{x} \in V$  and all  $y \in R$  with  $F(\bar{x}) \leq y < G(\bar{x})$ , (respectively  $F(\bar{x}) < y \leq G(\bar{x})$ ),  $(\bar{x}, y) \in U$ .

**Proof.** We prove the unparenthesized statement, the parenthetical one being similar.

Applying Lemma 3.1 we obtain an open cell  $D$  in  $R^n$ , with  $C \subseteq D$ , and a continuous definable retraction  $H : D \rightarrow C$ .

Let

$$V = \{\bar{x} \in D : d^{(n)}(\bar{x}, H(\bar{x})) < d^{(n+1)}((\bar{x}, f \circ H(\bar{x})), U^c)\},$$

where  $U^c = R^{n+1} \setminus U$ . Clearly  $V$  is open in  $R^n$  and (1) holds since  $\Gamma(f) \subseteq U$ . Putting  $F = f \circ H|_V$  we see that (2) holds. Also note that for all  $\bar{x} \in V$ ,  $F(\bar{x}) < g \circ H(\bar{x})$  and

$$J_{\bar{x}} := [0, g \circ H(\bar{x}) - F(\bar{x})] \subseteq \{t \in R_{\geq 0} : F(\bar{x}) + t \in U_{\bar{x}}\} \subseteq I$$

since  $U$  has  $I$ -short height.

By o-minimality and the fact that  $\Gamma(F) \subseteq U$ , there are well defined definable maps  $z_0 : V \rightarrow I$  and  $y_0 : V \rightarrow R$  given by

$$z_0(\bar{x}) = \sup\{t \in J_{\bar{x}} : [F(\bar{x}), F(\bar{x}) + t] \subseteq U_{\bar{x}}\}$$

and

$$y_0(\bar{x}) = F(\bar{x}) + z_0(\bar{x}).$$

Now observe that  $y_0 : V \rightarrow R$  satisfies the conditions (3), (4) and (5) for  $G$  ((3) is satisfied because  $(f, g)_C \subseteq U$ , by hypothesis, and  $f = F|_C$ ), but maybe  $y_0$  is not continuous. Thus we need to find a continuous definable map  $G : V \rightarrow R$  such that  $F < G \leq y_0$  and  $G|_C = y_0$ .

Consider the definable set

$$S = \{(\bar{x}, y) \in R^{n+1} : \bar{x} \in V \text{ and } F(\bar{x}) \leq y \leq g \circ H(\bar{x})\}$$

and the definable continuous maps  $\theta_1, \theta_2 : S \rightarrow I$  given by

$$\theta_1(\bar{x}, y) = 1_I -_I (y - F(\bar{x})) \cdot_I (g \circ H(\bar{x}) - F(\bar{x}))^{-1_I 1_I}$$

where  $1_I$  is the neutral element for the multiplication  $\cdot_I$ ,  $-_I$  is the difference and  $-^{1_I 1_I}$  is inversion in the field  $I$ , and,

$$\theta_2(\bar{x}, y) = \inf\{d^{(n+1)}((\bar{x}, t), U^c) : F(\bar{x}) \leq t \leq y\}.$$

Note that since  $U$  has  $I$ -short height we do have  $\theta_1(S) \subseteq I$  and  $\theta_2(S) \subseteq I$ .

Fix  $\bar{x} \in V$ . Then the continuous definable map  $(\theta_1 \cdot_I \theta_2)(\bar{x}, -)$  decreases monotonically and strictly from  $d^{(n+1)}((\bar{x}, F(\bar{x})), U^c)$  to  $0_I = 0$  on  $[F(\bar{x}), y_0(\bar{x})]$  and is identically  $0_I = 0$  on  $[y_0(\bar{x}), g \circ H(\bar{x})]$ .

For  $\bar{x} \in V$  let

$$a_{\bar{x}} = F(\bar{x}), \quad b_{\bar{x}} = y_0(\bar{x}), \quad c_{\bar{x}} = 0, \quad d_{\bar{x}} = d^{(n+1)}((\bar{x}, F(\bar{x})), U^c),$$

and  $\theta_{\bar{x}}(-) = (\theta_1 \cdot_I \theta_2)(\bar{x}, -)| : [a_{\bar{x}}, b_{\bar{x}}] \rightarrow [c_{\bar{x}}, d_{\bar{x}}]$ . Then by Remark 3.5,

$$G : V \rightarrow R : \bar{x} \mapsto \theta_{\bar{x}}^{-1}(d^{(n)}(\bar{x}, H(\bar{x})))$$

is a continuous definable function. Moreover,  $a_{\bar{x}} < G(\bar{x}) \leq b_{\bar{x}}$  for all  $\bar{x} \in V$ . In fact, if not then  $a_{\bar{x}} = G(\bar{x})$  and we obtain  $(\theta_1 \cdot_I \theta_2)(\bar{x}, G(\bar{x})) = d^{(n+1)}((\bar{x}, F(\bar{x})), U^c)$  contradicting the fact that  $d^{(n)}(\bar{x}, H(\bar{x})) < d^{(n+1)}((\bar{x}, F(\bar{x})), U^c)$ . We also have  $G(\bar{x}) = b_{\bar{x}}$  for all  $\bar{x} \in C$ . Therefore,  $G$  satisfies (3), (4) and (5) as required.

□

We need one more lemma:

**Lemma 3.7.** *Let  $C$  be a cell in  $R^n$ . Suppose that  $f, g : C \rightarrow R$  are continuous definable maps such that  $f < g$  and let  $U$  be an open definable subset of  $R^{n+1}$ . Suppose further that  $[f, g]_C \subseteq U$ . Then there exists an open definable subset  $W$  of  $R^n$  and continuous definable maps  $F, G : W \rightarrow R$  such that:*

- (1)  $C \subseteq W$ ;
- (2)  $F|_C = f$  and  $\Gamma(F) \subseteq U$ ;
- (3)  $G|_C = g$  and  $\Gamma(G) \subseteq U$ ;
- (4)  $F < G$ ;
- (5) for all  $\bar{x} \in W$  and all  $y \in R$  with  $F(\bar{x}) \leq y \leq G(\bar{x})$ ,  $(\bar{x}, y) \in U$ .

**Proof.** Applying Lemma 3.1 we obtain an open cell  $D$  in  $R^n$ , with  $C \subseteq D$ , and a continuous definable retraction  $H : D \rightarrow C$ .

Let  $W'$  be the intersection of

$$\{\bar{x} \in D : d^{(n)}(\bar{x}, H(\bar{x})) < d^{(n+1)}((\bar{x}, f \circ H(\bar{x})), U^c)\}$$

and

$$\{\bar{x} \in D : d^{(n)}(\bar{x}, H(\bar{x})) < d^{(n+1)}((\bar{x}, g \circ H(\bar{x})), U^c)\}$$

where  $U^c = R^{n+1} \setminus U$ . Clearly  $W'$  is open in  $R^n$  and (1) holds for  $W'$  since  $\Gamma(f), \Gamma(g) \subseteq U$ . Also (2) and (3) hold for  $f \circ H|_{W'}$  and  $g \circ H|_{W'}$ . Also note that for all  $\bar{x} \in W'$ ,  $f \circ H|_{W'}(\bar{x}) < g \circ H|_{W'}(\bar{x})$  so (4) holds for  $f \circ H|_{W'}$  and  $g \circ H|_{W'}$ .

Let  $B = [f \circ H|_{W'}, g \circ H|_{W'}]_{W'} \setminus U$  where

$$[f \circ H|_{W'}, g \circ H|_{W'}]_{W'} = \{(\bar{x}, y) \in W' \times R : y \in [f \circ H|_{W'}(\bar{x}), g \circ H|_{W'}(\bar{x})]\},$$

and let

$$W = W' \setminus \overline{\pi(B)}.$$

Clearly  $W$  is open. We now show that  $C \subseteq W$ , verifying in this way (1). Suppose not and let  $c = (c_1, \dots, c_n) \in C$  be such that  $c \in \pi(B)$ . Let  $\epsilon > 0$  be such that  $E = \prod_{i=1}^n [c_i - \epsilon, c_i + \epsilon] \subseteq W'$ . By definable choice there is a definable map  $\alpha : (0, \epsilon) \rightarrow \pi(B) \cap E$  such that  $\lim_{t \rightarrow 0^+} \alpha(t) = c$ . By replacing  $\epsilon$  we may assume that  $\alpha$  is continuous. Again by definable choice, we see that there exists a definable map  $\beta : (0, \epsilon) \rightarrow B \cap [f \circ H|_E, g \circ H|_E]_E$  such that  $\pi \circ \beta = \alpha$ . By replacing  $\epsilon$  we may assume that  $\beta$  is continuous. Since the definable set  $B \cap [f \circ H|_E, g \circ H|_E]_E$  is closed and, by [3, Proposition 3.1 (3)],  $\beta((0, \epsilon))$  is bounded, the limit  $\lim_{t \rightarrow 0^+} \beta(t)$  exists in this set. If  $d$  is this limit, then  $\pi(d) = c$  since  $\pi \circ \beta = \alpha$ . So  $d \in [f \circ H|_{W'}(c), g \circ H|_{W'}(c)] \cap B \neq \emptyset$  contradicting the fact that  $[f \circ H|_{W'}(c), g \circ H|_{W'}(c)] = [f(c), g(c)] \subseteq U$ .

If we put  $F = f \circ H|_W$  and  $G = g \circ H|_W$  we see that (2), (3) and (4) hold. On the other hand, if  $\bar{x} \in W$  and  $y \in R$  are such that  $F(\bar{x}) \leq y \leq G(\bar{x})$  and, by absurd,  $(\bar{x}, y) \notin U$ , then  $(\bar{x}, y) \in B$  and so  $\bar{x} \in \pi(B) \subseteq \overline{\pi(B)}$  contradicting the fact that  $\bar{x} \notin \overline{\pi(B)}$ . Thus (5) also holds. □

Combining Lemmas 3.6 and 3.7 we obtain:

**Lemma 3.8.** *Let  $C$  be a cell in  $R^n$ . Suppose that  $f, g : C \rightarrow R$  are continuous definable maps such that  $f < g$  and let  $U$  be an open definable subset of  $R^{n+1}$ . Suppose further that  $[f, g]_C \subseteq U$  (respectively  $(f, g)_C \subseteq U$ ). Then there exists a cell*

decomposition  $C_1, \dots, C_l$  of  $C$  and for each  $i = 1, \dots, l$  there is an open definable subset  $V_i$  of  $R^n$  and continuous definable maps  $F_i, G_i : V_i \rightarrow R$  such that:

- (1)  $C_i \subseteq V_i$ ;
- (2)  $F_i|_{C_i} = f|_{C_i}$  and  $\Gamma(F_i) \subseteq U$  (respectively  $\Gamma(G_i) \subseteq U$ );
- (3)  $G_i|_{C_i} = g|_{C_i}$ ;
- (4)  $F_i < G_i$ ;
- (5) for all  $\bar{x} \in V_i$  and all  $y \in R$  with  $F_i(\bar{x}) \leq y < G_i(\bar{x})$ , (respectively  $F_i(\bar{x}) < y \leq G_i(\bar{x})$ ),  $(\bar{x}, y) \in U$ .

**Proof.** We prove the unparenthesized statement, the parenthetical one being similar.

Let  $H : D \rightarrow C$  be as in Lemma 3.1. Choose  $\epsilon \in I$  such that  $2\epsilon \in I$  and put

$$U_f = U \cap ((f \circ H) - \epsilon, (f \circ H) + \epsilon)_D$$

and

$$U_g = U \cap ((g \circ H) - \epsilon, (g \circ H) + \epsilon)_D.$$

Then clearly  $U_f$  and  $U_g$  are open definable subsets of  $U$  with  $I$ -short height. For example, if  $(\bar{x}, y) \in U_f$ , then  $(f \circ H)(\bar{x}) - \epsilon < y < (f \circ H)(\bar{x}) + \epsilon$ .

Since  $(f, g)_C \subseteq U$ ,  $\Gamma(f) \subseteq U_f$  and  $\Gamma(g) \subseteq U_g$ , by Lemma 3.3, there exist definable maps  $f', g' : C \rightarrow R$  and cells  $C_1, \dots, C_m \subseteq C$  such that:

- (1)  $C = C_1 \cup \dots \cup C_m$ ;
- (2) for each  $i$ ,  $f'|_{C_i}$  and  $g'|_{C_i}$  are continuous;
- (3)  $f < f' < g' < g$ ;
- (4) for each  $i$ ,  $\Gamma(f'|_{C_i}) \subseteq U_f$  and  $\Gamma(g'|_{C_i}) \subseteq U_g$ ;
- (5) for each  $i$ ,  $(f'|_{C_i}, g|_{C_i})_{C_i} \subseteq U$ ,  $(f|_{C_i}, g'|_{C_i})_{C_i} \subseteq U$  and  $[f'|_{C_i}, g'|_{C_i}]_{C_i} \subseteq U$ .

Fix  $i = 1, \dots, m$ . Then we can apply Lemma 3.6 to the data  $(U_f, f|_{C_i}, f'|_{C_i})$  and obtain the data  $(V_f, F_1, F'_1)$  satisfying (1) to (5) of that lemma. Similarly, we can apply Lemma 3.6 to the data  $(U_g, g'|_{C_i}, g|_{C_i})$  and obtain the data  $(V_g, G'_1, G_1)$  satisfying (1) to (5) of that lemma. On the other hand, we can apply Lemma 3.7 to the data  $(U, f'|_{C_i}, g'|_{C_i})$  and obtain the data  $(W, F', G')$  satisfying (1) to (5) of that lemma.

Take  $V_i = V_f \cap V_g \cap W$  and set  $F = F_1|_{V_i}$ ,  $G = G_1|_{V_i}$ . Then clearly (1) to (5) hold.  $\square$

The following is also required:

**Lemma 3.9.** *Let  $C$  be a cell in  $R^n$ . Suppose that  $k : C \rightarrow R$  is a continuous definable map and let  $U$  be an open definable subset of  $R^{n+1}$ . Suppose further that  $[k, +\infty)_C \subseteq U$  (respectively  $(-\infty, k]_C \subseteq U$ ). Then there exists an open definable subset  $W$  of  $R^n$  and a continuous definable map  $K : W \rightarrow R$  such that:*

- (1)  $C \subseteq W$ ;
- (2)  $K|_C = k$  and  $\Gamma(K) \subseteq U$ ;
- (3) for all  $\bar{x} \in W$  and all  $y \in R$  with  $K(\bar{x}) \leq y$  (respectively  $y \leq K(\bar{x})$ ),  $(\bar{x}, y) \in U$ .

**Proof.** We prove the unparenthesized statement, the parenthetical one being similar.

Applying Lemma 3.1 we obtain an open cell  $D$  in  $R^n$ , with  $C \subseteq D$ , and a continuous definable retraction  $H : D \rightarrow C$ .

Let

$$W' = \{\bar{x} \in D : d^{(n)}(\bar{x}, H(\bar{x})) < d^{(n+1)}((\bar{x}, k \circ H(\bar{x})), U^c)\}$$

where  $U^c = R^{n+1} \setminus U$ . Clearly  $W'$  is open in  $R^n$  and (1) holds for  $W'$  since  $\Gamma(k) \subseteq U$ . Also (2) holds for  $k \circ H|_{W'}$ .

Let  $B = [k \circ H|_{W'}, +\infty)_{|W'} \setminus U$  where

$$[k \circ H|_{W'}, +\infty)_{|W'} = \{(\bar{x}, y) \in W' \times R : k \circ H|_{W'}(\bar{x}) \leq y\},$$

and let

$$W = W' \setminus \overline{\pi(B)}.$$

Clearly  $W$  is open. We now show that  $C \subseteq W$ , verifying in this way (1). Suppose not and let  $c = (c_1, \dots, c_n) \in C$  be such that  $c \in \overline{\pi(B)}$ . Let  $\epsilon > 0$  be such that  $E = \Pi_{i=1}^n [c_i - \epsilon, c_i + \epsilon] \subseteq W'$ . By definable choice there is a definable map  $\alpha : (0, \epsilon) \rightarrow \pi(B) \cap E$  such that  $\lim_{t \rightarrow 0^+} \alpha(t) = c$ . By replacing  $\epsilon$  we may assume that  $\alpha$  is continuous. Again by definable choice, we see that there exists a definable map  $\beta : (0, \epsilon) \rightarrow B \cap [k \circ H|_E, +\infty)_{|E}$  such that  $\pi \circ \beta = \alpha$ . By replacing  $\epsilon$  we may assume that  $\beta$  is continuous. Since the definable set  $B \cap [k \circ H|_E, +\infty)_{|E}$  is closed and, by [3, Proposition 3.1 (3)],  $\beta((0, \epsilon))$  is bounded, the limit  $\lim_{t \rightarrow 0^+} \beta(t)$  exists in this set. If  $d$  is this limit, then  $\pi(d) = c$  since  $\pi \circ \beta = \alpha$ . So  $d \in [k \circ H|_{W'}(c), +\infty) \cap B \neq \emptyset$  contradicting the fact that  $[k \circ H|_{W'}(c), +\infty) = [k(c), +\infty) \subseteq U$ .

If we put  $K = k \circ H|_W$  we see that (2) holds. On the other hand, if  $\bar{x} \in W$  and  $y \in R$  are such that  $K(\bar{x}) \leq y$  and, by absurd,  $(\bar{x}, y) \notin U$ , then  $(\bar{x}, y) \in B$  and so  $\bar{x} \in \pi(B) \subseteq \overline{\pi(B)}$  contradicting the fact that  $\bar{x} \notin \overline{\pi(B)}$ . Thus (3) also holds.  $\square$

**Corollary 3.10.** *If  $\mathcal{R}$  is a semi-bounded non-linear o-minimal expansion of an ordered group, then every non-empty open definable set is a finite union of open cells.*

**Proof.** This is done by induction on the dimension of the open definable set. For dimension one this is clear. Let  $U$  be an open definable subset of  $R^{n+1}$ . Let  $\mathcal{D}$  be a cell decomposition of  $R^{n+1}$  partitioning  $U$ . Clearly it is enough to show that each cell  $D \in \mathcal{D}$  with  $D \subseteq U$  can be covered by finitely many open cells (in  $R^{n+1}$ ) each of which is contained in  $U$ .

Case A:  $D = (f, g)_C$  for some cell  $C$  in  $R^n$  and continuous definable maps  $f, g : C \rightarrow R$  such that  $f < g$ .

Let  $f' = \frac{2f+g}{3}$  and  $g' = \frac{f+2g}{3}$ . Then  $f', g' : C \rightarrow R$  are continuous definable maps such that

- $f < f' < g' < g$ ;
- $\Gamma(f') \subseteq U$  and  $\Gamma(g') \subseteq U$ ;
- $(f', g')_C \subseteq U$  and  $(f, g')_C \subseteq U$ .

Now apply Lemma 3.8 to the data  $(C, U, f, g')$  and obtain the data  $(C_i, V_i, F_i, G'_i)$  with  $i = 1, \dots, l$  satisfying (1) to (5) of that lemma. By the inductive hypothesis there exists a finite collection  $\mathcal{A}_i$  of open cell in  $R^n$  contained in  $V_i$  which cover  $V_i$ . By (4) and (5) of Lemma 3.8, for each  $A \in \mathcal{A}_i$ ,  $(F_{i|A}, G'_{i|A})_A$  is an open cell in  $R^{n+1}$  contained in  $U$ , and by (1), (2) and (3) of that lemma,  $(f|_{C_i}, g'_{i|C_i})_{C_i} \subseteq \cup\{(F_{i|A}, G'_{i|A})_A : A \in \mathcal{A}_i\}$ . Thus  $(f, g')_C \subseteq \cup\{(F_{i|A}, G'_{i|A})_A : A \in \mathcal{A}_i \text{ and } i = 1, \dots, l\}$ .



Similarly, apply Lemma 3.8 to the data  $(C, U, f', g)$  (the parenthetical statement there) to see that  $(f', g)_C$  can be covered by finitely many open cells in  $R^{n+1}$  each of which is contained in  $U$ . Hence the same is true for  $(f, g)_C = (f, g')_C \cup (f', g)_C$ .

Case B:  $D = \Gamma(h)$  for some continuous definable map  $h : C \rightarrow R$  where  $C$  is a cell in  $R^n$ . This case reduces to Case A above by Lemma 3.2.

Case C:  $D = (k, +\infty)_C$  (respectively  $D = (-\infty, k)_C$ ) for some cell  $C$  in  $R^n$  and continuous definable map  $k : C \rightarrow R$ .

Then we can apply Lemma 3.9 to the data  $(C, U, k)$  and obtain the data  $(C, W, K)$  satisfying (1) to (3) of that lemma. By the inductive hypothesis there exists a finite collection  $\mathcal{A}$  of open cell in  $R^n$  contained in  $W$  which cover  $W$ . By (3) of Lemma 3.9, for each  $A \in \mathcal{A}$ ,  $(K|_A, +\infty)_A$  is an open cell in  $R^{n+1}$  contained in  $U$ , and by (1) and (2) of that lemma,  $(k|_C, +\infty)_C \subseteq \cup\{(K|_A, +\infty)_A : A \in \mathcal{A}\}$ .

Similarly for the case  $D = (-\infty, k)_C$ .  $\square$

## REFERENCES

- [1] S. Andrews *Definable open sets as finite unions of definable open cells* Notre Dame J. Formal Logic **51** (2010) 247–251
- [2] L. van den Dries *Tame Topology and o-minimal Structures* Cambridge University Press 1998.
- [3] M. Edmundo *Structure theorems for o-minimal expansions of groups* Ann. Pure Appl. Logic **102** (1-2) (2000) 159–181.
- [4] M. Edmundo and P. Eleftheriou *Definable group extensions in semi-bounded o-minimal structures* Math. Log. Quart. **55** (2009) 598–604.
- [5] M. Edmundo, P. Eleftheriou and L. Prelli *The universal covering map in o-minimal expansions of groups* Preprint (2011).
- [6] P. Eleftheriou *Local analysis for semi-bounded groups* Fund. Mathematicae **216** (2012) 223–258.
- [7] P. Eleftheriou and Y. Peterzil *Definable quotients of locally definable groups* Selecta Math. (N.S.) **18** (2012) 885–903.
- [8] P. Eleftheriou and Y. Peterzil *Definable groups as homomorphic images of semi-linear and field-definable groups* Selecta Math. (N.S.) **18** (4) (2012) 905–940.
- [9] P. Eleftheriou and S. Starchenko *Groups definable in ordered vector spaces over ordered division rings* J. Symb. Logic **72** (2007) 1108–1140.
- [10] J. Loveys and Y. Peterzil, *Linear o-minimal structures* Israel J. Math. **81** (1993) 1–30.
- [11] D. Marker, Y. Peterzil and A. Pillay *Additive reducts of real closed fields* J. Symb. Logic **57** (1992) 109–117.
- [12] Y. Peterzil *A structure theorem for semibounded sets in the reals* J. Symb. Logic **57** (1992) 779–794.
- [13] Y. Peterzil *Returning to semi-bounded sets* J. Symb. Logic **74** (2009) 597–617.
- [14] Y. Peterzil and S. Starchenko *A trichotomy theorem for o-minimal structures* Proc. London Math. Soc. **77** (3) (1998) 481–523.
- [15] A. Pillay, P. Scowcroft and C. Steinhorn *Between groups and rings* Rocky Mountain J. Math. **19** (3) (1989) 871–885.
- [16] A. Wilkie *Covering open definable sets by open cells* In: O-minimal Structures, Proceedings of the RAAG Summer School Lisbon 2003, Lecture Notes in Real Algebraic and Analytic Geometry (M. Edmundo, D. Richardson and A. Wilkie eds.), Cuvillier Verlag 2005.

UNIVERSIDADE ABERTA AND CMAF UNIVERSIDADE DE LISBOA, AV. PROF. GAMA PINTO 2,  
1649-003 LISBOA, PORTUGAL

*E-mail address:* `edmundo@cii.fc.ul.pt`

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO, 200 UNIVERSITY AVE WEST,  
N2L 3G1 WATERLOO, ONTARIO, CANADA

*E-mail address:* `pelefthe@uwaterloo.ca`

CMAF UNIVERSIDADE DE LISBOA, AV. PROF. GAMA PINTO 2, 1649-003 LISBOA, PORTUGAL

*E-mail address:* `lprelli@math.unipd.it`